

LOOK WHO'S BACK



METHAR® TRI-KOMBI

Herbicide



The Most Trusted Broadleaf Herbicide

Active Ingredients of

Methar Tri Kombi

336g/L Mecoprop,

80g/L 2,4-D

40g/L Dicamba

Available in 20L & 5L
Containers

GROUP

4

HERBICIDE



COLIN CAMPBELL (CHEMICALS) PTY LTD • (02) 9725 2544
HORTICULTURE • POST HARVEST • TURF

80 Years in 2020 • Est. 1940 • campbellchemicals.com.au
100% Australian owned & operated.



Methar Tri Kombi is registered for use on the following weeds at the rate of 4.2L/ha

Can be used at 3L/ha for newly sown turf (refer ot label)

Clovers	Chickweed	Bindii	Cobblers Peg
Creeping Oxalis	Lambs Tongue	Fumitory	Capeweed
Pearlwort	Catsear	Staggerweeds	Wireweed
Mouse-eared chickweed	Fat Hen	Mallow	

Best Use Practices

1. Ensure weeds are actively growing before herbicide application and turf is not mowed 2 days prior or after to ensure best possible results. This allows better absorption.
2. Allow at least 4-10 hours before irrigation or rain to ensure that the spray will be absorbed, the longer the better.
3. Avoid spraying in very hot dry conditions and if frost is likley to occur.
4. Using a suitable surfactant (such as Filmstar) on difficult weeds (must be used on oxalis). This will increase the distribution and adhesion of the spray hence increasing the efficacy.
5. The timing of sprays can be crucial. Weeds are always best sprayed when they are young; at this stage they are most susceptible to herbicides.
6. Spray until weeds and stem are just wet. Spraying beyond this point will encourage run off away from the target area with loss of efficacy.
7. Seeding can take place the day after. However do not apply to newly seeded turf for at least 4 weeks after first mowing (usually 8-12 weeks after germination). Then use the low rate of 30mL/100m².



@campbellchemicals



@campbellchemicals



Colin Campbell Chemicals



@campbellturf

**FilmStar - ideal to
add to our herbicide
products for better
sticking and spreading**

